

|             |   |
|-------------|---|
| Title       | Notes on carnivore fossils from the Pliocene Udunga fauna, Transbaikal area, Russia |
| Author(s)   | Ogino, Shintaro; Nakaya, Hideo; Takai, Masanaru; Fukuchi, Akira                     |
| Citation    | Asian paleoprimateology (2009), 5: 45-60  |
| Issue Date  | 2009  |
| URL         | <a href="http://hdl.handle.net/2433/199776">http://hdl.handle.net/2433/199776</a>   |
| Right       |   |
| Type        | Departmental Bulletin Paper   |
| Textversion | publisher   |

## Notes on carnivore fossils from the Pliocene Udunga fauna, Transbaikial area, Russia

Shintaro Ogino<sup>1\*</sup>, Hideo Nakaya<sup>2</sup>, Masanaru Takai<sup>1</sup>, Akira Fukuchi<sup>2, 3</sup>

<sup>1</sup>*Primate Research Institute, Kyoto University, Inuyama 484-8506, Japan*

<sup>2</sup>*Graduate School of Science and Engineering, Kagoshima University, Kagoshima  
890-0065, Japan*

<sup>3</sup>*Graduate School of Natural Science and Technology, Okayama University, Okayama  
700-8530, Japan*

*\*Corresponding author. e-mail: ogino@pri.kyoto-u.ac.jp*


### Abstract

We provide notes of carnivore fossils from the middle Pliocene Udunga fauna, Transbaikial area, Russia. The fossil carnivore assemblage consists of more than 200 specimens including eleven genera. *Ursus*, *Parailurus*, *Parameles*, and *Ferienstrix* are representative of the animals of thermophilic forest biotopes. On the other hand, *Chasmaporthetes* and *Pliocrocota* are probably specialized in open environment. The prosperity both in forestal and semiarid carnivores indicate that the Udunga fauna is comprised of mosaic elements.

### Introduction

The fauna dating back to the Pliocene in Udunga, Transbaikalia, Russia, comprises eleven species of mammals, such as rodents, lagomorphs, carnivores, perissodactyls, artiodactyls, and elephants (Kalmykov, 1989, 1992, 2003; Kalmykov and Maschenko, 1992, 1995; Vislobokova *et al.*, 1993, 1995; Erbajeva *et al.*, 2003). The Udunga site is located on the left bank of the Temnik River, the tributary of the Selenga River in the vicinity of Udunga village (Figure 1). This mammal fauna is discovered from the red clayey beds of the Upper Neogene Chikoi Formation (Kalmykov, 1989; Erbajeva *et al.*, 2003).

The fossil materials of the Pliocene Transbaikalian carnivores have been studied by many workers (e.g. Kalmykov, 1989, 1992, 2003; Sotnikova and Kalmykov, 1991; Vislobokova *et al.*, 2001; Sotnikova, 2006, 2008), and these series are their ongoing research. Their works show the rich diversity of carnivores in the Pliocene Transbaikalian area and the richness is particular feature of the Udunga fauna. In this study, we remark eleven genera of carnivores from the Udunga locality, which have been recognized from the materials stored in the Southern Scientific Center (SSC), Rostov-on-Don, Russia. The preliminary study of carnivorous mammals of Udunga fauna was reported by Sotnikova and Kalmykov (1991),



**Figure 1.** The location of the Transbaikal area. Modified from Kalmykov (2002).

but *Canis* sp., *Gulo minor* are not stored in SSC collection. The recognized species in this study are listed in Table 1.

### Remarks

#### **Order Carnivora Bowdich, 1821**

#### **Family Canidae Gray, 1821**

#### **Genus *Nyctereutes* Temmnick, 1838**

#### ***Nyctereutes* sp. indet.**

Sotnikova and Kalmykov (1991) reported the occurrence of *Nyctereutes* from the Udunga fauna. Three hemimaxillae, four hemimandibles, isolated  $M^2$ , and three of isolated  $M_1$  of *Nyctereutes* are distinguished from the SSC collection (Table 2). There are a few *Nyctereutes* materials known in the Udunga fauna, which is identified as *Nyctereutes* sp. (Sotnikova and Kalmykov, 1991; Kalmykov, 1992, 1999, 2003; Erbajeva *et al.*, 2003; Kalmykov, Maschenko, 2005, 2006; Kalmykov *et al.*, 2005; Maschenko *et al.*, 2007). The oldest representative of the genus *Nyctereutes* (*N. aff. donnezani*) is known from late Turolian Venta del Moro (Spain), so the origin is considered to appear in western part of Palaearctic region (Morales and, Aquirre, 1976). In Ruscinian, they were widespread in Europe (Alcalá and Montoya, 1989-1990; Tedford, 1995). Early representatives of *Nyctereutes* (*N. donnezani*) are known from La Gloria 4 (MN 14), La Calera (MN 15a), Layna (MN 15b),

**Table 1.** List of carnivore fossils of the Udunga fauna stored in SSC collection.

| taxon                           | number of specimens | references                    |
|---------------------------------|---------------------|-------------------------------|
| <i>Nyctereutes</i> sp. indet    | 11                  | Sotnikova and Kalmykov (1991) |
| <i>Canis</i> sp. indet.         | 2                   | Sotnikova and Kalmykov (1991) |
| <i>Ursus minimus</i>            | 8                   | Sotnikova and Kalmykov (1991) |
| <i>Pannonictis</i> sp. indet.   | 8                   | Vislobokova et al (2001)      |
| <i>Parameles suillus</i>        | 5                   | Kalmykov (1989)               |
| <i>Ferinestris</i> sp. indet.   | 107                 | Sotnikova (2006)              |
| <i>Parailurus baicalicus</i>    | 3                   | Sotnikova (2008)              |
| <i>Pliocrocota pyrenaica</i>    | 30                  | Kalmykov (1989)               |
| <i>Chasmaporthetes lunensis</i> | 4                   | Vislobokova et al (2001)      |
| <i>Lynx shansius</i>            | 28                  | Kalmykov (1989)               |
| <i>Homotherium</i> sp. indet.   | 2                   | Kalmykov (1989)               |

**Table 2.** Specimens of *Nyctereutes* sp. indet.

| Udg- | side | Material  |
|------|------|---|
| 43   | R    | maxilla with M <sup>1</sup>                                 |
| 48   | R    | mandible with M <sub>1</sub> -M <sub>3</sub>                |
| 49   | L    | mandible with P <sub>4</sub> and M <sub>1</sub>             |
| 50   | R    | maxilla with P <sup>4</sup> -M <sup>2</sup>                 |
| 119  | R    | mandible with lower canine, P <sub>2</sub> , P <sub>4</sub> |
| 161  | R    | maxilla with P <sub>3</sub> and P <sub>4</sub>              |
| 162  | R    | M <sub>1</sub> (bloken)                                     |
| 163  | L    | M <sub>1</sub>  |
| 190  | L    | M <sup>2</sup>  |
| ##   | R    | mandible with M <sub>1</sub>                                |
| ##   | R    | M <sub>1</sub>  |

**Table 3.** Specimens of *Canis* sp. indet.

| Udg- | side | Material                     |
|------|------|------------------------------|
| 176  | L    | mandible with M <sub>2</sub> |
| 2471 |      | upper and lower dentitions   |

Spain. Later form, *N. megamastoides* is from Villarroja (MN 16a ), El Rincon (MN 16b), La Puebla del Valverde (MN 17a), Spain (Alcalá and Montoya, 1989-1990). *Nyctereutes* is also known from the western Transbaikalia area (Sotnikova and Kalmykov, 1991, Kalmykov, 1992) and China (Tedford, Qiu, 1991; Tong *et al.*, 1995; Li *et al.*, 2003) in Villafranchian. In Central to East Asia, *N. cf. sinensis* is known from western Transbaikalia (Beregovaya), and *N. megamastoides* is known from the northern Mongolia (Shamar) (Sotnikova, 1976). Tedford and Qui (1991) hypothesized that the early Pliocene Chinese and European *Nyctereutes* should be organized in the same taxon. *N. megamastoides* diverged from *N. donnezani* at the late Pliocene in Europe (Soria and Aguirre, 1976), which is morphologically comparable with *N. sinensis*. On the other hand, *N. tingi*, which is known from East Asia and suddenly occurred in the early Pliocene, is closely related to extant *N. sinensis* (Tedford and Qui, 1991). Thus, identification of *Nyctereutes* from the Udunga fauna would be important for understanding relationship between European and Asian taxon in Pliocene.

### Genus *Canis* Linnaeus, 1758

#### *Canis* sp. indet.

Only two specimens of large canid are available in the SSC Udunga collection (Table 3). Left mandible with M<sub>2</sub> (Udg-0176) and isolated upper and lower dentitions (Udg-2471)

**Table 4.** Specimens of *Ursus* sp. indet.

| Udg- | side | Material                    | Udg- | side | Material       |
|------|------|-----------------------------|------|------|----------------|
| 16   | R    | lower canine                | 2001 |      | upper canine   |
| 53   | L    | M <sub>2</sub>              | 2002 | L    | M <sub>1</sub> |
| 54   | R    | M <sub>2</sub>              | 2003 | R    | M <sub>2</sub> |
| 140  | L    | maxilla with P <sup>4</sup> | 2004 | R    | M <sub>3</sub> |

**Table 5.** Specimens of *Pannonictis* sp. indet.

| Udg- | side | Material                                       |
|------|------|--|
| 128  | L    | maxilla with P <sup>4</sup>                    |
| 129  | R    | maxilla with P <sup>3</sup> and P <sup>4</sup> |
| 144  | R    | P <sup>4</sup>                                 |
| 145  | R    | M <sup>1</sup>                                 |
| 150  | L    | P <sup>4</sup>                                 |
| 2400 | R    | upper canine                                   |
| 2410 | L    | upper canine                                   |
| 2400 | L    | lower canine                                   |

are obviously larger than *Nyctereutes*. These specimens are considered *Canis* but there are some small forms. *Canis* appeared in Eurasia in the middle Turolian (Mein, 1990), but their adaptive radiation occurred in Ruscinian (Vangengeim *et al.*, 1998). Its oldest representatives are known from Europe such as Los Mansuetos, Concud (*C. cipio*, MN 12), Perpingnan (*C. michauxi*, *C. adoxus*, MN 15), La Calera (*C. indeterminado*, MN 15a), Odessa catacombs (*C. petenyi*, MN 15), Villarroja (*C. indeterminado*, MN 16a), Kuruksai (*C. kuruksaensis*, *C. ex gr. lephagus*, MN 16b), Kvabebi (*Canis* sp., MN 16), Etouaires (*C. etruscus*, MN 16b) (Vekua, 1972; Vangengeim *et al.*, 1988; Alcalá and Montoya, 1989-1990; Mein, 1990). In China, *Canis* appeared in the late Ruscinian (Tedford, 1995; Tedford, Qiu, 1996).

### Family Ursidae Fisher de Waldheim, 1817

#### Genus *Ursus* Linnaeus, 1758

#### *Ursus minimus* Devèze et Bouillet, 1827 (Stuart, 1982)

Sotnikova and Kalmykov (1991), Kalmykov (1992) reported *Ursus ex gr. ruscinensis-minimus* from the Udunga fauna. Seven isolated teeth including upper canine, lower canine, M<sub>1</sub>, M<sub>2</sub>, M<sub>3</sub>, and a left maxilla with P<sup>4</sup>, are stored in the SSC Udunga collection (Table 4). *U. minimus* retains primitive morphological characteristics, which is closely related to the late Ruscinian or the earliest Villafranchian ursid forms (Sotnikova and Kalmykov, 1991; Kalmykov, 1992; Sotnikova, 2006). *U. minimus* Deveze et Debouillet (sin. *U. arvernensis* Croizet et Jobert) is known from three locality, such as Gaville (Val de'Arno, Italy), Puy de Dôme (France) and Layna (Spain) (Torres, 1988).

The identification of the Pliocene Russian ursid is disputed (Vekua, 1972; Sotnikova and Kalmykov, 1991; Kalmykov, 1992; Vislobokova *et al.*, 1993; and Sotnikova, 2006). In Europe *Ursus* is known from the late Ruscinian localities, such as Layna (*U. ruscinensis*), Odessa catacombs (*U. arvernensis*), Weze (*U. wenzensis*) (Kowalski, 1990; Mein, 1990; Perez, Soria, 1989-1990; Vangengeim *et al.*, 1998). In China *Ursus* sp. is known from Ertente (*Ursus* sp.). The stratigraphical range of genus *Ursus* is from the late Turolian to the present.

### Family Mustelidae Fischer de Waldheim, 1817

**Table 6.** Specimens of *Parameles* sp. indet.

| Udg- | side | Material  |
|------|------|---|
| 2    | L    | mandible with lower canine-M <sub>2</sub>                   |
| 3    | L    | mandible with lower canine-M <sub>2</sub>                   |
| 177  | L    | mandible with lower canine, P <sub>4</sub> , M <sub>1</sub> |
| 178  | L    | mandible with M <sub>1</sub> and M <sub>2</sub>             |
| 2420 |      | upper canine  |

**Subfamily Galictinae Reig, 1957****Genus *Pannonictis* Kormos, 1931*****Pannonictis* sp. indet.**

Vislobokova *et al.* (2001) reported the occurrence of large galictine mustelid, *Pannonictis*, from Asian part of former USSR (Table 5). Eight of mustelid specimens including six isolated tooth, left maxilla with P<sup>4</sup>, and right maxilla with P<sup>3</sup> and P<sup>4</sup>, considered to be *Pannonictis*. *Pannonictis* differs from other Mustelidae in the sizes and shape of M<sub>1</sub> and P<sup>4</sup>, skull, and mandible. It is close to the South American extant genus *Grison* and to the extinct genus *Enhydriactis* from the Plio-Pleistocene of Europe (Sotnikova, 1980). The Middle Pleistocene East Asian *Oriensictis* is also close to *Pannonictis* (Ogino and Otsuka, 2008). The origin of *Pannonictis* is probably in Asia. Zdansky (1927) has described a large specimen of Mustelidae gen. et sp. indet. from the accumulations of late Miocene of China, which was similar to *Pannonictis*. Sotnikova (1980) considered that both of *Pannonictis* from the northern Mongolia (Shamar, MN 16a) and China (Gaotege, MN 15) are relatives of the Zdansky's mustelid. In Europe the oldest *Pannonictis* is known from the late Ruscinian of Hungary (Csarnota 1, MN 15) (García and Howell, 2008).

**Subfamily Melinae Burmeister, 1850****Genus *Parameles* Rostshin, 1949*****Parameles suillus* (Teilhard et Leroy) 1945**

Kalmykov (1989) reported the occurrence of *Parameles suillus* from the Udunga fauna. Four left mandibles and isolated upper canine are considered to be *Parameles suillus* in the SSC Udunga collection (Table 6). Left mandible (Udg-0003) is described by Kalmykov (2003: p. 31, Fig.4). *P. suillus* (= *Meles suillus*), which has been described from adjournment of the sediments of Yushe (Teilhard de Chardin and Leroy, 1945), was found from Shamar (northern Mongolia) (Sotnikova, 1980). Several melid fossils have already been described under the different names from the early Pliocene Europe: *Parameles* from Odessa catacombs (Ukraine) and Weze (Poland) (Stach, 1951). According to Sotnikova and Kalmykov (1991), the priority belongs to *Parameles* Rostchin, so *Parameles* should include “*Meles*” *genevauxi* from Montopoli (France), *Arctomeles pliocaenicus* from Weze (Poland), *Parameles ferus* from the Odessa catacombs (Ukraine), *Meles suillus* from Yushe, Ertemte (China), Shamar (North

**Table 7.** Specimens of *Ferinetrix* sp. indet.

| Udg- | side | Material   | Udg- | side | Material  |
|------|------|--|------|------|---|
| 4    | R    | mandible with M <sub>1</sub>                                       | 167  | R    | mandible with lower canine and P <sub>2</sub>   |
| 6    | L    | mandible with P <sub>3</sub> , P <sub>4</sub> , and M <sub>2</sub> | 168  | R    | mandible with M <sub>1</sub>                    |
| 7    | L    | mandible with lower canine-M <sub>2</sub>                          | 169  |      | upper incisor                                   |
| 8    | L    | mandible with P <sub>3</sub> -M <sub>1</sub>                       | 170  | L    | mandible with P <sub>2</sub> -M <sub>1</sub>    |
| 9    | R    | mandible with P <sub>3</sub> -M <sub>1</sub>                       | 171  | R    | mandible with P <sub>3</sub> -M <sub>2</sub>    |
| 10   | L    | mandible with I <sub>2</sub> , lower canine-M <sub>1</sub>         | 172  | L    | mandible with P <sub>4</sub> -M <sub>2</sub>    |
| 11   | L    | mandible with P <sub>3</sub>                                       | 173  | L    | mandible with lower canine-M <sub>1</sub>       |
| 12   | R    | mandible with P <sub>3</sub>                                       | 174  | R    | mandible with P <sub>2</sub> -M <sub>1</sub>    |
| 13   | L    | mandible with lower canine-M <sub>2</sub>                          | 175  | R    | maxilla with P <sup>3</sup> -M <sup>1</sup>     |
| 14   | L    | mandible with P <sub>4</sub> and M <sub>1</sub>                    | 185  | L    | mandible with lower canine-M <sub>1</sub>       |
| 18   | L    | mandible with P <sub>2</sub> and P <sub>4</sub>                    | 1297 | R    | mandible with P <sub>4</sub> and M <sub>1</sub> |
| 19   | L    | maxilla with P <sup>4</sup>  | 1302 | L    | upper canine                                    |
| 20   | L    | mandible with P <sub>4</sub> -M <sub>1</sub>                       | 1303 | R    | M <sup>1</sup>                                  |
| 31   | L    | mandible with P <sub>3</sub> -M <sub>1</sub>                       | 1304 | L    | P <sup>3</sup>                                  |
| 32   | L    | mandible with m2   | 2005 | R    | lower canine                                    |
| 46   | L    | mandible with P <sub>4</sub> -M <sub>2</sub>                       | 2006 | L    | maxilla with P <sup>3</sup> and P <sup>4</sup>  |
| 47   | R    | mandible with lower canine-M <sub>2</sub>                          | 2007 | L    | mandible with M <sub>1</sub>                    |
| 55   | L    | M <sub>1</sub>   | 2009 | R    | upper canine                                    |
| 56   | R    | M <sub>1</sub>   | 2010 | L    | upper canine                                    |
| 57   |      | lower canine   | 2011 | R    | M <sup>1</sup>                                  |
| 68   | L    | M <sub>1</sub>   | 2012 | R    | M <sub>2</sub>                                  |
| 69   | L    | maxilla with P <sup>3</sup> and P <sup>4</sup>                     | 2013 | L    | M <sub>2</sub>                                  |
| 70   | L    | M <sup>1</sup>   | 2016 | R    | maxilla with M <sup>1</sup>                     |
| 72   | L    | mandible with lower canine-M <sub>1</sub>                          | 2017 | R    | maxilla with P <sup>4</sup> and M <sup>1</sup>  |
| 73   | L    | mandible with lower canine-M <sub>2</sub>                          | 2018 | L    | maxilla with M <sup>1</sup>                     |
| 74   | L    | mandible with M <sub>1</sub> and M <sub>2</sub>                    | 2031 | R    | mandible  |
| 75   | R    | maxilla with P <sup>4</sup> and M <sup>1</sup>                     | 2005 | L    | lower canine                                    |
| 79   | L    | maxilla with P <sup>4</sup> and M <sup>1</sup>                     | 2016 | L    | M <sup>1</sup>                                  |
| 82   | L    | mandible with P <sub>4</sub> and M <sub>1</sub>                    | 2017 | R    | M <sup>1</sup>                                  |
| 83   | L    | maxilla with P <sup>4</sup> and M <sup>1</sup>                     | 2348 |      | upper canine                                    |
| 89   | R    | mandible with lower canine-M <sub>1</sub>                          | 2349 |      | upper canine                                    |
| 90   | L    | M <sup>1</sup>   | 2352 |      | upper canine                                    |
| 91   | R    | mandible with P <sub>4</sub> and M <sub>1</sub>                    | 2353 |      | upper canine                                    |
| 111  | R    | mandible with M <sub>1</sub>                                       | 2459 |      | upper canine                                    |
| 113  | R    | mandible with P <sub>4</sub> and M <sub>1</sub>                    | 2369 |      | upper canine                                    |
| 114  | L    | mandible with M <sub>1</sub>                                       | 2370 |      | upper canine                                    |
| 115  | L    | mandible with M <sub>1</sub>                                       | 2410 |      | lower canine                                    |
| 116  | L    | mandible with P <sub>3</sub> -M <sub>1</sub>                       | 2411 |      | lower canine                                    |
| 117  | L    | mandible with P <sub>3</sub> -M <sub>1</sub>                       | 2412 |      | lower canine                                    |
| 122  | R    | M <sup>1</sup>   | 2413 |      | lower canine                                    |
| 123  | L    | maxilla with P <sup>3</sup> and P <sup>4</sup>                     | 2415 |      | upper canine                                    |
| 124  | R    | P <sup>4</sup>   | 2416 |      | upper canine                                    |
| 125  | R    | M <sup>1</sup>   | 2439 |      | upper canine                                    |
| 130  | R    | maxilla with P <sup>3</sup>  | 2431 |      | upper canine                                    |
| 132  | R    | mandible with P <sub>3</sub> -M <sub>1</sub>                       | 2428 |      | upper canine                                    |
| 133  | R    | mandible with lower canine, P <sub>3</sub> , P <sub>4</sub>        | 2429 |      | upper canine                                    |
| 135  | R    | maxilla with P <sup>4</sup> and M <sup>1</sup>                     | 2424 |      | lower canine                                    |
| 136  | L    | M <sup>1</sup>   | 2425 |      | lower canine                                    |
| 137  | L    | maxilla with P <sup>4</sup> and M <sup>1</sup>                     | 2426 |      | lower canine                                    |
| 141  | L    | maxilla with P <sup>4</sup>  | 2454 |      | lower canine                                    |
| 158  | R    | maxilla with P <sup>3</sup> and P <sup>4</sup>                     | 2455 |      | lower canine                                    |
| 159  | L    | maxilla with P <sup>3</sup> - M <sup>1</sup>                       | 2472 |      | upper canine                                    |
| 164  | L    | maxilla with M <sup>1</sup>  | 2408 | R    | M <sub>1</sub>                                  |
| 165  | R    | maxilla with P <sup>4</sup> and M <sup>1</sup>                     |      |      |   |

Mongolia) and Udunga. Thus, the stratigraphical range of *P. suillus* is considered from the end of Turolian (MN 13) to the beginning of early Villafranchian (MN 16a) (Li *et al.*, 1984; Wolsan, 1989; Sotnikova, Kalmykov, 1991; Kalmykov, 1992; Tedford, 1995; Vangengeim *et al.*, 1998).

**Table 8.** Specimens of *Parailurus baikalicus*.

| Udg- | side | Material                                  |
|------|------|---|
| 1    | R    | mandible with lower canine-M <sub>2</sub> |
| 51   | R    | M <sub>2</sub>                            |
| 52   | R    | M <sub>1</sub>                            |

**Genus *Ferinestrix* Bjork, 1970*****Ferinestrix* sp. indet.**

An enigmatic form of large badger is identified as *Ferinestrix* sp. by Sotnikova (2006). Formerly a large badger was defined as Mustelidae gen. nov. et sp. nov (Kalmykov, 1989). The most abundant carnivore in the SSC Udunga collection is *Ferinestrix* (Table 7), which was only known from the Pliocene North American Hagerman local fauna and established on the basis of the fragment of mandible with incomplete M<sub>1</sub> (Bjork, 1970).

**Family Ailuridae Gray, 1843****Genus *Parailurus* Schlosser, 1899*****Parailurus baikalicus* Sotnikova, 2008**

Only three specimens are available to us (Table 8). Right mandible with complete cheek teeth row (Udg-0001) has been reported by Kalmykov (2002: p. 211, Fig. 3). All dentition of Udg-0001 are considerably worn, indicating that it belong to an old individual, while two isolated teeth, right M<sub>1</sub> (Udg-0052) and M<sub>2</sub> (Udg-0051), have weak dental attrition.

These ailurid specimens are considered to belong to the large lesser panda, *Parailurus baikalicus* Sotnikova, 2008 from Udunga. Although extant lesser panda, *Ailurus fulgens* is limited to the Himalayan highland forest, the extinct large lesser panda *Parailurus* is widely distributed in the Pliocene northern hemisphere including Europe, Asia, and North America (Kundrát, 1997; Morlo and Kundrát, 2001; Sasagawa *et al.*, 2003; Wallace and Wang, 2004; Sotnikova, 2008). Several specimens of Ailuridae are known from the Udunga fauna and numerous lists of fauna was specified as *Ailurus* sp. (Kalmykov, 1999, 2003, Erbajeva *et al.*, 2003; Kalmykov, Maschenko, 2005, 2006; Kalmykov *et al.*, 2005; Maschenko *et al.*, 2007). Erbajeva *et al.* (2003) identified these specimens to *Parailurus* sp., but later it was redescribed as *Parailurus baikalicus* (Sotnikova, 2008) on the basis of the fragment left maxilla (No. 962/58, БФ ГИИ) with P<sup>4</sup>, M<sup>1-2</sup> and the alveolus of the distal root of P<sup>3</sup>. The Transbaikalian *P. baikalicus* is larger than the European *P. anglicus*, but likely corresponds to *P. hungaricus*. *Parailurus* sp. from the Pliocene Japan (Sasagawa *et al.*, 2003) differs from *P. baikalicus* by having primitive characteristics on P<sup>4</sup>.



**Table 9.** Specimens of *Pliocrocota* sp. indet.

| Udg- | side | Material  | Udg- | side | Material                                       |
|------|------|---|------|------|--|
| 24   | R    | mandible with P <sub>2</sub> -M <sub>1</sub>                | 187  | L    | maxilla with P <sup>3</sup> and P <sup>4</sup> |
| 25   | R    | mandible with lower canine, P <sub>4</sub> , M <sub>1</sub> | 188  | L    | maxilla with P <sup>2</sup> -P <sup>4</sup>    |
| 26   | R    | mandible with lower canine-M <sub>1</sub>                   | 1301 | L    | I <sub>1</sub> -I <sub>3</sub>                 |
| 27   | L    | mandible with P <sub>2</sub> -M <sub>1</sub>                | 2008 | L    | P <sup>3</sup>                                 |
| 29   | R    | mandible with lower canine-M <sub>1</sub>                   | 2014 | R    | M <sub>1</sub>                                 |
| 30   | L    | mandible with M <sub>1</sub>                                | 2015 | R    | mandible                                       |
| 41   | R    | mandible with P <sub>2</sub> -M <sub>1</sub>                | 2019 | R    | mandible                                       |
| 71   | R    | mandible with lower canine-M <sub>1</sub>                   | 2020 |      | lower incisor                                  |
| 100  | L    | P <sup>4</sup>  | 2021 | R    | I <sub>3</sub>                                 |
| 156  | R    | mandible with P <sub>3</sub> and P <sub>4</sub>             | 2022 | L    | I <sub>3</sub>                                 |
| 180  | R    | mandible with P <sub>2</sub> -M <sub>1</sub>                | 2028 | R    | mandible with P <sub>3</sub> -M <sub>1</sub>   |
| 181  | R    | mandible with P <sub>3</sub> -M <sub>1</sub>                | 2032 | R    | M <sub>1</sub>                                 |
| 182  | L    | mandible with lower canine-M <sub>1</sub>                   | 2033 | L    | P <sub>3</sub>                                 |
| 183  | R    | mandible with lower canine-M <sub>1</sub>                   | 2034 | L    | P <sub>4</sub>                                 |
| 186  | R    | maxilla with P <sup>3</sup> and P <sup>4</sup>              | 2035 | L    | M <sub>1</sub>                                 |

**Family Hyaenidae Gray, 1869****Genus *Pliocrocota* Koretzoï, 1938*****Pliocrocota pyrenaica* (Deperet), 1890**

30 specimens are considered to be *Pliocrocota pyrenaica* (Table 9). Right mandible (Udg-0026) of *P. pyrenaica* has been reported by Kalmykov (2003: p. 32, Fig.5). *P. pyrenaica* retains primitive morphological feature and similar to the late Ruscinian or the earliest Villafranchian hyaenid form (Kalmykov, 1992; Sotnikova, 2006). The identification of *Pliocrocota* is constructed based on tooth morphology. The characteristics of M<sub>1</sub> differs from that of *Hyaena*. The length of P<sub>4</sub> is much greater than that of *Ictitherium* (Sotnikova, 1989). *Pliocrocota* possibly includes fossil hyenids from Layna (*Hyaena pyrenaica*), Odessa catacombs (*Crocota* cf. *sivalensis*), Beregovaya (*Crocota* cf. *licenti*), Kuruksai (*Pachycrocota* cf. *perrieri*), Leijiahe (*Hyaena licenti*) (Kalmykov, personal communication). *P. pyrenaica* is common not only in Udunga but also in the early Pliocene of Europe, such as Serrat den Vocquer (France), Layna (Spain), and Odessa catacombs (Ukraine). *Pliocrocota* is distributed in the wide area of northern Eurasian continent from Spain to the Baikal lake in the early Pliocene (Vangengeim *et al.*, 1998). In the Pliocene of Europe some stages of the development of the genus from *P. pyrenaica* of Ruscinian to the derived form, *P. perrieri* (Sotnikova, 1989). In Central Asia *Pliocrocota* is known from the early to middle Pleistocene. Kurten (1956) marked finds of two large forms such as *P. brevirostris licenti* and *P. brevirostris sinensis* in Zhoukoudian, northern China. Subspecies of *P. brevirostris* is also known from Transbaikalia (Zasukhina, Tologoi) and from the Northern Mongolia (Nalaiha) (Vislobokova *et al.*, 1995). *P. perrieri* is reported from the late Pliocene localities in Asia, such as Youhe (China) and Kuruksay (southern Tajikistan) (Howell, Pether, 1980; Sotnikova, 1989). In Asia, *Pachycrocota pyrenaica orientalis* is known from the Villafranchian China (Qiu, 1987). Unfortunately, the Chinese material on hyaenid is not stratified, therefore new data on *P. pyrenaica* from the western Transbaikalia area would be

**Table 10.** Specimens of *Chasmaporthetes* sp. indet.

| Udg- | side | Material                                     |
|------|------|--|
| 45   | L    | mandible with P <sub>2</sub> -M <sub>1</sub> |
| 184  | L    | mandible with P <sub>2</sub> -M <sub>1</sub> |

**Table 11.** Specimens of *Lynx shansius*.

| Udg- | side | Material  | Udg- | side | Material  |
|------|------|---|------|------|---|
| 15   | R    | mandible with lower canine-M <sub>1</sub>                   | 127  | R    | maxilla with P <sup>3</sup> and P <sup>4</sup>              |
| 22   | R    | mandible with lower canine-M <sub>1</sub>                   | 134  | R    | mandible with P <sub>3</sub> and P <sub>4</sub>             |
| 33   | L    | mandible with lower canine, P <sub>4</sub> , M <sub>1</sub> | 138  | L    | mandible with lower canine, P <sub>3</sub> , P <sub>4</sub> |
| 34   | R    | mandible with lower canine, P <sub>4</sub> , M <sub>1</sub> | 139  | R    | mandible with P <sub>4</sub> and M <sub>1</sub>             |
| 35   | R    | mandible with lower canine-M <sub>1</sub>                   | 157  | L    | mandible with I <sub>3</sub> and P <sub>3</sub>             |
| 36   | R    | mandible with lower canine-M <sub>1</sub>                   | 160  | R    | mandible with lower canine-M <sub>1</sub>                   |
| 38   | R    | mandible with P <sub>3</sub> -M <sub>1</sub>                | 166  | L    | maxilla with P <sup>4</sup> and M <sup>1</sup>              |
| 39   | L    | mandible with P <sub>3</sub> -M <sub>1</sub>                | 179  | L    | mandible with lower canine-P <sub>4</sub>                   |
| 40   | L    | mandible with lower canine-M <sub>1</sub>                   | 1295 | L    | mandible with P <sub>3</sub> -M <sub>1</sub>                |
| 44   | L    | mandible with P <sub>3</sub> -M <sub>1</sub>                | 1305 |      | lower canine  |
| 77   | R    | maxilla with P <sup>3</sup> and P <sup>4</sup>              | 1306 |      | lower canine  |
| 86   | R    | mandible with lower canine-M <sub>1</sub>                   | 1307 |      | lower canine  |
| 110  | R    | mandible with P <sub>3</sub> -M <sub>1</sub>                | 1308 | R    | mandible with lower canine and P <sub>3</sub>               |
| 118  | L    | mandible with P <sub>3</sub> and P <sub>4</sub>             | 1309 | R    | P <sub>4</sub>  |

important for the study of its dispersal and evolution.

### Genus *Chasmaporthetes* Hay, 1921

#### *Chasmaporthetes* sp. indet.

Vislobokova *et al.* (2001) reported the occurrence of a gracile hunting hyaena, *Chasmaporthetes*, from Asian part of the former USSR. Only two left mandibles with P<sub>2</sub>-M<sub>1</sub> considered to be this genus (Table 10). *Chasmaporthetes* was widely distributed in the Plio-Pleistocene northern hemisphere including North Africa, Europe, Asia, and North America (e.g. Hay, 1921). In Asia *Chasmaporthetes* (*C. lunensis*) is discovered from the late Pliocene of the Northern Mongolia (Shamar) and Udunga. The fossil specimens of the hunting hyaena are discovered in western Transbaikalia area (Beregovaya) and Tajikistan (Kuruskay), and have been described as *Euryboas* (Sotnikova, 1974, 1976, Vangengeim *et al.*, 1988). Repenning (1967) has proposed an assumption that *Euryboas* is a synonym of genus *Chasmaporthetes*. Savage and Curtis (1970) and Kurtén and Werdelin (1988) supported his opinion. However, the priority of the genus *Chasmaporthetes* has been restored even later (Kurten and Crusafont, 1977). According to Sotnikova (1994), *Chasmaporthetes* includes *C. borissiakii* (presumably late Miocene, Moldova), *C. exitelus* (late Miocene, China), *C. lunensis* (Pliocene - early Pleistocene, Eurasia), *C. ossifragus* (late Pliocene - early Pleistocene, North America), *C. nitidula* (late Pliocene - early Pleistocene, Africa). Thus, chronological distribution of *Chasmaporthetes* is from the late Miocene to the early Pleistocene.

**Table 12.** Specimens of *Homotherium* sp. indet.

| Udg- | side | Material  |
|------|------|---|
| 21   | R    | mandible with lower canine, P <sub>4</sub> , M <sub>1</sub> |
| 23   | R    | mandible with P <sub>4</sub> and M <sub>1</sub>             |
| 76   | R    | P <sup>4</sup>  |
| 101  | R    | P <sup>4</sup>  |

**Family Felidae Gray, 1821****Genus *Lynx* Kerr, 1792*****Lynx shansius* Teilhard et Leroy, 1945**

Kalmykov (1989) reported the occurrence of *Lynx shansius* from the Udunga locality. 28 specimens are considered to be this species in the SSC Udunga collection (Table 11) and a right mandible (Udg-0160) has been described by Kalmykov (2003: p. 34, Fig. 6). *Lynx* from Udunga is known to be one of the oldest lynxes: its morphology is more primitive than that of *Lynx shansius* from the Shamar and Yushe and *L. issiodorensis* from Pardines (France) and Val d'Arno (Italy) (Kurten and Werdelin, 1984). *Lynx* appeared in the late Ruscinian and is known from Moldova and Odessa catacombs (Vangengeim *et al.*, 1998).

**Genus *Homotherium* Fabrini, 1890*****Homotherium* sp. indet.**

Kalmykov (1989) firstly reported the occurrence of *Homotherium* sp. from the Udunga locality. Two right mandibles and two isolated right P<sup>4</sup> are considered to be *Homotherium* in the SSC Udunga collection (Table 12). *Homotherium* is considered to have separated from *Machairodus* at the end Miocene (Anton and Galobart, 1999), but *H. crenatidens* is known from the Villafranchian of Eurasia (Bajgusheva, 2000; Hemmer, 2001; Sotnikova *et al.*, 2002). The saber-toothed cat, Machairodontinae, is discovered from the Ruscinian locality of Europe and Asia, but the materials are so fragmentary to define their essential characteristics (Vekua, 1972; Perez, Soria, 1989-1990). *Homotherium* sp. from the Ruscinian Odessa catacombs (Ukraine), differs from the Villafranchian specimen in its primitive characteristics (Vangengeim *et al.*, 1998).

**Discussion**

Eleven genera of carnivores have been recognized from the SSC Udunga collection (Table 1). The Udunga fauna is recognized as the end of early Pliocene (Kalmykov, 1989, 1992, 2003; Sotnikova and Kalmykov, 1991; Kalmykov and Malaeva, 1994; Kalmykov and Maschenko, 2005, 2006; Kalmykov *et al.*, 2005; Maschenko *et al.*, 2007; Maschenko *et al.*, 2007) or the early Late Pliocene (MN 16a, Vislobokova *et al.*, 1993; Vislobokova *et al.*, 1995; Erbajeva *et al.*, 2003) on the basis of biostratigraphical study of many kind of mammals.

However, the Pliocene Eurasian carnivores were widely distributed spatiotemporally, and carnivore taxa of the Udunga fauna have close relationships not only to the Eurasian but also to the African and North American assemblages.

Vislobokova *et al.* (1995) referred that the occurrence of *Parameles suillus* and *Ursus minimus* from the Udunga fauna indicates analogues with Beregovaya assemblage (MN 16b; Transbaikal, Russia) and Shamar assemblage (2.8 Ma; Mongolia). The former carnivoran assemblage is represented by *Nyctereutes sinensis*, *Eucyon cf. minimus*, *Chasmaporthetes lunensis*, *Lynx shansius* and *Acinonyx cf. pardinensis* (Alexeeva and Erbajeva, 2006). The latter carnivoran assemblage is represented by *Nyctereutes megamostoides*, *Mustela* sp., *Pannonictis pachygnata*, *Meles (=Parameles) suillis*, *Hyaena cf. licenti*, *Euriboas cf. lunensis*, *Felis (=Lynx) shansius*, and *Acinonyx* sp. (Sotnikova, 1980 and Zazhigin, 1989). Vislobokova *et al.* (1995) also pointed out that *Ursus minimus*, *Pachycrocuta (=Pliocrocuta) pyrenaica*, and felids are comparable to the Italian Triversa fauna (MN 16a; Azzaroli *et al.*, 1988). The Khapry faunal assemblage reported from the coastal zone of Azov Sea, Russia, (MN 17) was recently revised by Sotnikova *et al.* (2002), where the carnivoran assemblage is represented by *Nyctereutes megamostoides*, *Canis cf. senecensis*, *Lutra* sp., *Pannonictis nesti*, *Pliocrocuta perrieri*, *Pachycrocuta brevirostris*, *Homotherium crenatidens*, *Acinonyx pardinensis*, and *Lynx issiodorensis*. Sotnikova (2008) referred that the similarity of *Pliocrocuta* and *Parameles* from the Udunga fauna to corresponding forms from the fauna of Odessa Catacombs, Ukraine (terminal Ruscinian).

Qiu and Qiu (1995) summarized the Chinese Neogene mammalian faunas and described the two Pliocene faunas from Shanxi Province. In the Yushe basin, *Pliohyaena (=Pliocrocuta) pyrenaica*, *Chasmaporthetes*, and *Nyctereutes* have been discovered from Gaozhuang fauna with a paleomagnetic age of 5.2 - 3.4 Ma. Deng *et al.* (2004) reported the Early Pliocene (MN14 - MN15) Hewangjia mammalian assemblage, Gansu Province, which includes three carnivoran genera, *Promephitis* sp., *Chasmaporthetes* sp., and *Hyaenictitherium wongi*, while *Ursus*, *Nyctereutes*, and *Canis* have not been found there. The absence of these forestal elements indicates that the sedimentary paleoenvironment of the Hewangjia fauna was dry condition.

*Ursus*, *Parameles*, and *Pliocrocuta* of the Udunga fauna show primitive features (Sotnikova, 2006) and these taxon borrow(???) heavily from the Ruscinian form. *Chasmaporthetes*, *Parailurus*, *Gulo*, and felids are discovered across the Northern Hemisphere including Europe, Asia, North Africa, and North America. The enigmatic large badger, *Ferinstrix*, was only known from the Pliocene North American Hagerman local fauna (Bjork, 1970). As mentioned above, carnivores of the Udunga fauna are considerable to the evidence of active exchanges between the New World and Old World in the Pliocene and this fauna is important for discussing about mammalian dispersal event in the Northern Hemisphere at the period. In addition, this fact shows the difficulty to use carnivore to the biostratigraphical study.

On the other hand, feeding habitat of carnivore is clearly discriminated from the fossil materials and it is useful to reconstruct the paleoenvironment. The abundance of animals of thermophilic forest biotopes and their unusual composition (with *Ursus*, *Parailurus*, *Parameles*, and *Ferienstrix*) is peculiar feature of carnivores of the Udunga fauna (Sotnikova, 2006). However, *Hipparion* (*H. houfenese* and *H. tchicoicum*), *Dicerorhinus*, bovids, and *Chasmaporthetes* are representative of the open land elements. The occurrence of both forestal and semiarid mammals indicates that the Udunga fauna is comprised of mosaic elements.

### Acknowledgements

We are deeply grateful to Prof. N.P. Kalmykov of the Southern Scientific Center, Russian Academy of Sciences, and Dr. E.N. Maschenko of the Paleontological Institute, Russian Academy of Sciences, for providing us with useful the carnivoran fossil specimens of the Udunga fauna. We are grateful to Dr. Marina V. Sotnikova (Geological Institute, Russian Academy of Sciences) for providing us with useful information on carnivora assemblage of the Udunga fauna and suggesting pertinent literature. This research was supported by JSPS (Japan Society for the Promotion of Science) and RFBR (Russian Foundation for Basic Research) under the Japan-Russia Research Cooperative Project (too M. Takai, Grant 840064200002 and to E.N. Maschenko, Grant 060491469).

### References

- Alcalá, L. and Montoya P. (1989-1990) Las faunas de macromamíferos del Turoliense inferior español, *Paleontologia i Evolució* **23**: 111–119.
- Alexeeva, N.V., Erbajeva M. A. (eds.) (2006) Stratigraphy, paleontology and paleoenvironment of Pliocene – Pleistocene of Transbaikalia and interregional correlations. Field Excursion Guidebook. INQUA. 14–23.
- Antón, M. and Galobart A. (1999) Neck function and predatory behavior in the scimitartoothed cat *Homotherium latidens* (Owen). *Journal of the Vertebrate Paleontology* **19** (4): 771–784.
- Azzaroli, A., De Giuli, C., Ficarelli, G., and Torre, G. (1988) Late Pliocene to early mid-Pleistocene mammals in Eurasia: faunal successions and dispersal events. *Palaeogeographia Italica* **71**: 48–54.
- Bajgusheva, V.S. (2000) New data on the Tamanian fauna complex from the section near Semibalki village (the Sea of Azov region). In: Kiyashko, V. (Ed.), *Historical–archeological Investigations in Azov and on the Lower Don River in 1998*. (Azov): 27–57. [in Russian].
- Bjork, P.R. (1970) The Carnivora of the Hagerman Local Fauna (Late Pliocene) of Southwestern Idaho. *Transactions of the American Philosophical Society* **60**(7): 1–54.
- Deng, T., Wang, X., Ni, X., Liu, L., and Liang, Z. (2004) Cenozoic stratigraphic sequence of the Linxia Basin in Gansu, China and its evidence from mammal fossils. *Vertebrata Palasiatica* **42**(1): 41–66.
- Erbajeva, M., Alexeeva, N., and Khenzykhenova, F. (2003) Pliocene small mammals from the Udunga site of the Transbaikal area. *Coloquios de Paleontologia* **1**: 133–145.
- Fabrini, A. (1890) *Machairodus* (Meganthereon) del Val d’Arno superiore. *Bollettino del Reale comitato geologico d’Italia*. ser. 3, v. 1: 121–144, 161–177.
- Ficarelli, G. (1979) The Villafranchian machairodonts of Tuscany. *Palaeontographia Italica* **71**: 17–26.

- García, N., Howell, F.C. (2008) New discovery of a large mustelid (Carnivora: Mammalia) from the early Pleistocene locality of Sima del Elefante (Sierra de Atapuerca, Spain). *Palaeontographica Abteilung A: Palaeozoologie - Stratigraphie* **284**: 1-16.
- Hay, O.P. (1921) Descriptions of species of Pleistocene vertebrata, types or specimens most of which are preserved in the United States National Museum. *Proceedings of the United States National Museum* 59: 599–642.
- Hemmer, H. (2001) Die Feliden aus dem Epivillafranchium von Untermassfeld. In: Kahlke, R.D (ed.) *Das Pleistozän von Untermassfeld bei Meiningen (Thüringen) Teil 3. Monographien des Römisch-Germanisches Zentralmuseums*. Bonn, vol. 40: 699–782.
- Howell, F.C. and Petter, G. (1989) The *Pachycrocuta* and *Hyaena* lineages (Plio-Pleistocene and extant species of the Hyaenidae). Their relationships with Miocene Ictitheres: *Palhyaena* and *Hyaenictitherium*, *Geobios* 13: 579-623.
- Kalmykov, N.P. (1989) New location of the mammals' hipparion fauna in the Western Transbaikalia. *Bull. of Commission for study of the Quaternary* 58: 133-138. [in Russian]
- Kalmykov, N.P. (1992) Biostratigraphy and Fauna of the Transbaikalian Pliocene mammals. Nauka Press, Novosibirsk, 97 p. [in Russian].
- Kalmykov, N.P. (1999) *Ecosystem Evolution of the Lake Baikal Basin in the Late Cenozoic*. Buryat Scientific Centre RAS, Ulan-Ude (1999): 114p.
- Kalmykov, N.P. (2002) Prehistoric animals at Lake Baikal. Rostov Universitetata, Izdatelstvo Rostovskogo universiteta, Rostov-on-Don: 229 p. [in Russian]
- Kalmykov, N.P. (2003) ПАЛЕОГЕОГРАФИЯ И ЭВОЛЮЦИЯ БИОЦЕНОТИЧЕСКОГО ПОКРОВА В БАСЕЙНЕ ОЗЕРА БАЙКАЛ: 238 p.
- Kalmykov, N.P. and Malaeva, E.M. (1994) Continental Biota from the Lower Pliocene of the Western Transbaikal Region,” *Dokl. Akad. Nauk* 339 (6): 785–788.
- Kalmykov, N.P. and Maschenko E.N. (1992) The northernmost Early Pliocene Cercopithecidae from Asia. *Paleontological Journal* 26(2): 178-181.
- Kalmykov, N.P. and Maschenko E.N. (1995) New cercopithecoid genus (Cercopithecidae, Primates) from Pliocene of Transbaikalian. *Voprosy Antropologii* 82: 106-121 [in Russian].
- Kalmykov, N.P., Maschenko E.N., Takai M., Taru H., Hasegawa Y., Koizumi A., and Iwamoto M. (2005) Zoogeography of monkeys (Mammalia, Colobidae) and mammalian fauna from Central Asia in the Pliocene. In: *Evolution of Life on the Earth*. TSU Publishing House, Tomsk: 290-292. [in Russian]
- Kalmykov, N.P., Shabunova, V. V. (2005). The first find of the Asian raccoons (Ailurinae: Carnivora, Mammalia) in Russia. In: *Evolution of Life on the Earth*. TSU Publishing House, Tomsk: 293-295. [in Russian]
- Kowalski K. (1990) Stratigraphy of Neogene mammals in Poland. In: *E.H. Lindsay, V. Fahlbusch and P. Mein, eds., European Neogene Mammal Chronology*, Plenum Press, New York: 193-209.
- Kundrát M. (1997) New dental remains of an Extinct lesser panda- Morphotype or new species? *The Fifth International Congress of Vertebrate Morphology abstracts*: p. 58A.
- Kurtén, B. (1956) The status and affinities of *Hyaena sinensis* Owen and *Hyaena ultima* Matsumoto. *American Museum Novitates* 1764: 1-48.
- Kurtén, B. and Crusafont, P.M. (1977) Villafranchian Carnivores (Mammalia) from La Puebla de Valverde (Teruel, Spain). *Comm. Biolo.*, 85: 1-39.
- Kurtén, B. and Werdelin, L. (1988), B. and Werdelin L. (1984) The relationships of *Lynx shansius* Teilhard. *Annales Zoologici Fennici* 21: 129-133.
- Kurtén, B. and Werdelin, L. (1988) A review of the genus *Chasmaporthetes* Hay, 1921 (Carnivora,

- Hyaenidae). *Journal of the Vertebrate Paleontology* 8: 46–66.
- Li, C., Wu, W., Qiu Z. (1984) Chinese Neogene: subdivision and correlation. *Vertebrata Palasiatica* 22(3): 163-178.
- Li, Q., Wang, X.-M., and Qiu Z.-D. (2003) Pliocene mammalian fauna of Gaotege in Nei Mongol (Inner Mongolia), China. *Vertebrata Palasiatica*. 41: 104-114.
- Maschenko, E.N. 2005 Cenozoic primates of eastern Eurasia (Russia and adjacent areas). *Anthropological Science* 113: 103-115.
- Maschenko, E.N. 2006
- Maschenko, E.N., Takai, M., Kalmykov, N.P. (2007) Ecology of fossil Cercopithecoidea (Primate, Mammalia) from mountain territories of Asia. In: *Mammals of mountain territories*. Moscou: Association of scientific editions KMK: 203-208. [in Russian]
- Mazo and Torres (1989-1990)
- Mein P. (1990) Updating of MN zones. In: *E.H. Lindsay, V. Fahlbusch and P. Mein, (Eds) European Neogene Mammal Chronology*. Plenum Press, New York, London: 73–90.
- Morales J. and Aquirre E. (1976) Carnívoros de Venta del Moro, *Trabajos sobre Neógeno–Cuaternario* 5: 31–81.
- Morlo, M. and Kundrát M. (2001) The first carnivoran fauna from the Ruscium (Early Pliocene, MN 15) of Germany. *Palaeontologische Zeitschrift* 75(2): 163–187.
- Ogino, S. and Otsuka, H. (2008) New Middle Pleistocene Galictini (Mustelidae, Carnivora) from the Matsugae Cave Deposits, northern Kyushu, west Japan. *Paleontological Research*. 12(2): 159-166.
- Perez, B. and Soria, D. (1989-1990) Análisis de las comunidades de mamíferos del Plioceno de Layna (Soria) y La Calera (Teruel). *Paleontología i Evolució*, 23: 231-238.
- Qiu, Z. (1987) Die Hyaeniden aus dem Ruscium und Villafranchium Chinas, *Münchener Geowissenschaftliche Abhandlungen* 9: 1–110.
- Qiu, Z. and Qiu, Z. (1995) Chronological sequence and subdivision of Chinese Neogene mammalian faunas. *Palaeogeography, Palaeoclimatology, Palaeoecology* 116: 41–70.
- Repenning, C. A. (1967) Palearctic-Nearctic mammalian dispersals in the late Cenozoic. In: *Hopkins D. M. (Ed), The Bering land bridge*. Stanford University Press: 288–311.
- Sasagawa, I., Takahashi, K., Sakumoto, T., Nagamori, H., Yabe, H., and Kobayashi, I. (2003) Discovery of the extinct red Pand Parailurus (Mammalia, Carnivora) in Japan. *Journal of Vertebrate Paleontology* 23(4): 895–900.
- Savage, D.E. and Curtis, G.H. (1970) The Villafranchian Stage-Age and its Radiometric Dating. *Geological Society of America, Special Paper* 124: 207-231.
- Soria, D. and Aguirre, E. (1976) El cánido de Layna: Revisión de los *Nyctereutes* fósiles, *Trabajos Sobre Neógeno-Cuaternario* 5: 82–115.
- Sotnikova, M.V. (1976) The upper-Pliocene Carnivora of Central Asia. *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya*, 11: 133–137. [in Russian]
- Sotnikova, M.V. (1980) The late-Pliocene Mustelidae from the locality Shamar (Mongolian People's Republic). *Bulleten Komissii po Izucheniyu Chetvertichnogo Perioda*, 50: 138–145. [in Russian]
- Sotnikova, M.V. (1994) The genus *Chasmaporthetes* Hay, 1921 from the Pliocene of Russia, Ukraine, Mongolia and Tadzhikistan In: Tatarinov L.P. (ed.), *Palaeotheriology*. Nauka, Moscow: 113–139. (In Russian)
- Sotnikova, M.V. and Kalmykov, N.P. (1991) Pliocene carnivora from Udunga locality (Transbaikalia, USSR). In: *Vangengeim E.A. (Ed.) Pliocene and Anthropogene Paleogeography and Biostratigraphy, Geological Institute of Russian Academy of Sciences*. Nauka, Moscow: 146-160 [in Russian].

- Sotnikova, M.V., Baigusheva, V.S., and Titov, V.V. (2002) Carnivores of the Khapr Faunal Assemblage and Their Stratigraphic Implications *Stratigraphy and Geological Correlation* 10 (4): 375–390.
- Sotnikova, M.V. (2003) Review of the Olyorian carnivore assemblage in Northern Siberia. *Impacts of late Quaternary Climate change on western Arctic shelf-lands: Insights from the Terrestrial Mammal Record*, a workshop convened at The International Arctic Research Center, University of Alaska Fairbanks, Fairbanks, Alaska.
- Sotnikova, M.V. (2006) Pliocene-Early Pleistocene carnivore assemblages of Transbaikalian area, Russia. *Stratigraphy, Paleontology, and Paleoenvironment of Pliocene-Pleistocene of Transbaikalia and interregional correlations*. 84–85.
- Sotnikova, M.V. (2008) A New Species of Lesser Panda *Parailurus* (Mammalia, Carnivora) from the Pliocene Transbaikalia (Russia) and Some Aspects of Ailurine Phylogeny *Paleontological Journal* 42(1): 90–99.
- Stach, J. (1951) *Arctomeles pliocaenicus* new genus and species from Weze. *Acta Geologica Polonica* 2: 129–157.
- Stuart, A. J. (1982) Pleistocene vertebrates of the British Isles. Longman, London; New York: 212 p.
- Tedford, R.H. (1995) Neogene mammalian biostratigraphy in China: past, present and future. *Vertebrate Palasiatica* 33 (4), 277–289.
- Tedford, R.H. and Qiu, Z. (1991) Pliocene *Nyctereutes* (Carnivora, canidae) from Yushe, Shanxi, with comments on Chinese fossil raccoon-dogs, *Vertebrata Palasiatica* 29: 176–189.
- Tedford, R.H. and Qiu, Z. (1996) A new canid genus from the Pliocene of Yushe, Shanxi Province. *Vertebrata Palasiatica* 34: 27–40.
- Teilhard de Chardin, P. and Leroy, P. (1945) Les mustelidés de Chine. *Publications de l'Institut de Géobiologie, Pékin*. 50 p.
- Tong, Y., Zheng, S.H., and Qiu, Z.D. (1995) Cenozoic mammal ages of China. *Vertebrata Palasiatica* 33: 290–314.
- Torres, T (1988) Osos (Mammalia, Carnivora, Ursidae) del Pleistoceno Ibe'rico (*Ursus spelaeus* Ros.-Hein, *Ursus deningeri* Von. Reich. y *Ursus arctos* L.). I. Filogenia, distribución estratigráfica y geográfica. Estudio anatómico y métrico del cráneo. *Boletín Geológico y Minero* 99: 3–46.
- Vangengeim, E.A., Vislobokova, I.A., and Sotnikova, M.V., (1998) Large Ruscinian Mammalia in the territory of the former Soviet Union. *Stratigraphy. Geological Correlation* 6 (4): 368–382.
- Vekua, A.K. (1972). The Kvabebi fauna of the Akchagylian. *Vertebrates Science*, Moscow. 350 p. [in Russian with English abstract]
- Vislobokova, I. A., Erbajeva, M.V., and Sotnikova, M. A. (1993) The early Villafranchian stage in the development of the mammalian fauna of Northern Eurasia. *Stratigraphy, Geological Correlation* 1(5): 87–96. [in Russian]
- Vislobokova, I., Dmitrieva, E., and Kalmykov, N.P. (1995) Artiodactyls from the Late Pliocene of Udunga, Western Trans-Baikal, Russia *Journal of Vertebrate Paleontology* 15(1): 146–159.
- Vislobokova, I., Sotnikova, M., and Dodonov, A. (2001) Late Miocene – Pliocene mammalian faunas of Russia and neighbouring countries. *Bollettino della Società Paleontologica Italiana* 40(2): 307–313.
- Wallace, S.C. and Wang, X. (2004) Two new carnivores from an unusual late Tertiary forest biota in eastern North America. *Nature* 431(30): 556–559.
- Wolsan, M. (1989) Carnivora. In: Historia i ewolucja lądowej fauny Polski (K. Kowalski, ed.). *Folia Quaternaria* 59–60: 177–196 [in Polish].
- Zazhigin, V.S. (1989) Late Cenozoic of Mongolia. Zaitsev N.S., Krashenninnikov V.A., Gerbova V.G., Timofeev P.P., and Tomortogoo O. (Eds). 200 p.
- Zdansky, O. (1927) Weitere Bemerkungen fiber fossile Carnivoren aus China. *Palaeontologia Sinica*. Ser.



C. 3, Fasc. 4: 1-28.